



Scientific Inquiry and Review (SIR)

Volume 3, Issue 2, April 2019

ISSN (P): 2521-2427, ISSN (E): 2521-2435

Journal DOI: <https://doi.org/10.32350/sir>

Issue DOI: <https://doi.org/10.32350/sir.32>

Homepage: <https://ssc.umt.edu.pk/sir/Home.aspx>

Journal QR Code:



Article

On Leap Reduced Reciprocal Randic and Leap Reduced Second Zagreb Indices of Some Graphs

Fazal Dayan

Muhammad Javaid

Muhammad Aziz ur Rehman

Author(s)

Online
Published

April 2019

Article DOI

<https://doi.org/10.32350/sir.32.04>

Article QR
Code



Fazal Dayan

Dayan F, Javaid M, ur Rehman MA. On leap reduced reciprocal randic and leap reduced second Zagreb indices of some graphs. *Sci Inquiry Rev.* 2019;3(2):27–35.

[Crossref](https://doi.org/10.32350/sir.32.04)

To cite this
article

Copyright
Information

This article is open access and is distributed under the terms of Creative Commons Attribution – Share Alike 4.0 International License.

Indexing Agency



A publication of the
School of Science, University of Management and Technology
Lahore, Pakistan.

On Leap Reduced Reciprocal Randic and Leap Reduced Second Zagreb Indices of Some Graphs

Fazal Dayan^{1*}, Muhammad Javaid¹, Muhammad Aziz ur Rehman¹

¹Department of Mathematics, School of Science,
University of Management and Technology, Lahore, Pakistan

[*fazaldayan1@gmail.com](mailto:fazaldayan1@gmail.com)

Abstract

Naji et al. introduced the leap Zagreb indices of a graph in 2017 which are new distance-degree-based topological indices conceived depending on the second degree of vertices. In this paper, we have defined the first and second leap reduced reciprocal Randic index and leap reduced second Zagreb index for selected wheel related graphs.

Keywords: leap indices, reduced reciprocal Randic index, reduced second Zagreb index, flower graph, gear graph, helm graph, sunflower graph, wheel graph

1. Introduction

In the current chemical and mathematical literature, a large number of vertex degree based graph invariants have been studied. Among them, the first and second Zagreb indices are, by far, the most extensively investigated. They were introduced more than forty years ago [1].

The properties of the two Zagreb indices can be seen in [2, 3, 4, 5, 6]. Many novel variants of the Zagreb indices have been studied in recent years. Some of these are multiplicative Zagreb indices [7, 8, 6]. Zagreb coincides [9, 10], multiplicative Zagreb coincides [11] and the sum Zagreb index [12, 6]. Recently, leap Zagreb indices of a graph have been introduced by Naji et al. [13], which are new distance-degree-based topological indices conceived depending on the second degree of vertices (number of their second neighbors). Some basic properties of these new indices have been established as well. A. M. Naji and N. D. Soner [14] presented the exact expressions for the first Zagreb index of selected graph operations containing the corona product, cartesian product, composition, dis junction and symmetric difference between graphs. Shiladhar P. et al. [15] computed leap Zagreb indices of selected wheel related graphs. Fazal Dayan et al. [16] defined leap Gourava indices and computed the exact values of leap Gourava indices for some wheel related graphs. In recent years, many researchers have worked on computing topological indices [17, 18, 19].

2. Definitions and Preliminaries

Definition. The reduced reciprocal Randic index is defined as follows:-

$$RRR(G) = \sum_{uv \in E} \sqrt{(d(u)-1)(d(v)-1)}$$

Definition. The reduced second Zagreb index is defined as follows:-

$$RM_2(G) = \sum_{uv \in E} (d(u)-1)(d(v)-1)$$

Definition. The leap reduced reciprocal Randic and leap reduced second Zagreb indices are defined as follows:-

$$LRRR(G) = \sum_{uv \in E} \sqrt{(d_2(u)-1)(d_2(v)-1)} \quad (1)$$

and

$$LRM_2(G) = \sum_{uv \in E} (d_2(u)-1)(d_2(v)-1) \quad (2)$$




Figure 1. Wheel graph W_n




Figure 2. Gear graph G_n




Figure 3. Helm graph H_n




Figure 4. Flower graph Fl_n




Figure 5. Sunflower graph Sf_n

Figures 1-5 show the wheel graph, gear graph, helm graph, flower graph and sunflower graph, respectively.

3. Main Results

Theorem

Let W_n be a wheel graph with $n + 1$ vertices, its leap reduced reciprocal Randic and leap reduced second Zagreb indices are given below:-

$$\text{LRRR}(G) = n[\sqrt{4-n} + n(n-4)]$$

$$\text{LRM}_2(G) = n(n-4)(n-5)$$

Proof: We compute the required result by using definitions 1 and 2 given in the previous section as follows:-

$$\begin{aligned} \text{LRRR}(G) &= \sum_{uv \in G} \sqrt{(d_2(u)-1)(d_2(u)-1)} \\ &= n\sqrt{[(0-1)(n-3-1)]} + n\sqrt{[(n-3-1)(n-3-1)]} \\ &= n\sqrt{[(-1)(n-4)]} + n\sqrt{[(n-4)]^2} \\ &= n\sqrt{4-n} + n(n-4) \\ &= n[\sqrt{4-n} + n(n-4)] \end{aligned}$$

and

$$\begin{aligned} \text{LRM}_2(G) &= \sum_{uv \in G} (d_2(u)-1)(d_2(u)-1) \\ \text{LRM}_2(G) &= n[(0-1)(n-3-1)] + n[(n-3-1)(n-3-1)] \\ &= n(4-n) + n(4-n)^2 \\ &= n(4-n + n^2 - 8n + 16) \\ &= n(n^2 - 9n + 20) \\ &= n(n-4)(n-5) \end{aligned}$$

Theorem

Let G_n be a gear graph with $2n + 1$ vertices, its leap reduced reciprocal Randic and leap reduced second Zagreb indices are given below:-

$$\text{LRRR}(G) = n(n-2)[\sqrt{n-1} + 2\sqrt{2}]$$

$$\text{LRM}_2(G) = n(n-2)(5n-1)$$

Proof: We compute the required result by using definitions 1 and 2 given in the previous section as follows:-

$$\begin{aligned}
\text{LRRR}(G) &= \sum_{uv \in G} \sqrt{(d_2(u)-1)(d_2(u)-1)} \\
&= n\sqrt{[(n-1)(n-1-1)]} + 2n\sqrt{[(n-1-1)(3-1)]} \\
&= n\sqrt{[(n-1)(n-2)]} + 2n\sqrt{2(n-2)} \\
&= n(n-2)[\sqrt{n-1} + 2\sqrt{2}]
\end{aligned}$$

and

$$\begin{aligned}
\text{LRM}_2(G) &= \sum_{uv \in G} (d_2(u)-1)(d_2(u)-1) \\
&= n[(n-1)(n-1-1)] + 2n[(n-1-1)(3-1)] \\
&= n[(n-1)(n-2)] + 2n[2(n-2)] \\
&= n(n-2)[n-1 + 4n] \\
&= n(n-2)(5n-1)
\end{aligned}$$

Theorem

Let H_n be a helm graph with $2n + 1$ vertices, its leap reduced reciprocal Randic and leap reduced second Zagreb indices are below:-

$$\begin{aligned}
\text{LRRR}(G) &= n(n-2) [\sqrt{n-1} + \sqrt{(n-2)} + \sqrt{2}] \\
\text{LRM}_2(G) &= n(n-2)(2n-1)
\end{aligned}$$

Proof: We compute the required result by using definitions 1 and 2 given in the previous section as follows:-

$$\begin{aligned}
\text{LRRR}(G) &= \sum_{uv \in G} \sqrt{(d_2(u)-1)(d_2(u)-1)} \\
&= n\sqrt{[(n-1)(n-1-1)]} + n\sqrt{[(n-1-1)(n-1-1)]} + n\sqrt{[(n-1-1)(3-1)]} \\
&= n\sqrt{[(n-1)(n-2)]} + n\sqrt{(n-2)^2} + n\sqrt{2(n-2)} \\
&= n(n-2) [\sqrt{n-1} + \sqrt{(n-2)} + \sqrt{2}]
\end{aligned}$$

and

$$\begin{aligned}
\text{LRM}_2(G) &= \sum_{uv \in G} (d_2(u)-1)(d_2(u)-1) \\
&= n[(n-1)(n-1-1)] + n[(n-1-1)(n-1-1)] + n[(n-1-1)(3-1)] \\
&= n[(n-1)(n-2)] + n[(n-2)(n-2)] + 2n[(n-2)] \\
&= n(n-2)(n-1 + n-2 + 2) \\
&= n(n-2)(2n-1)
\end{aligned}$$

Theorem

Let F_n be a flower graph with $2n + 1$ vertices, its leap reduced reciprocal Randic and leap reduced second Zagreb indices are given below:-

$$\begin{aligned} \text{LRRR}(G) &= n \left[\sqrt{[(2n-3)(2n-5)]} + \sqrt{3-2n} + \sqrt{5-2n} + (2n-5) \right] \\ \text{LRM}_2(G) &= 8n(n^2-5n+6) \end{aligned}$$

Proof: We compute the required result by using definitions 1 and 2 given in the previous section as follows:-

$$\begin{aligned} \text{LRRR}(G) &= \sum_{uv \in E} \sqrt{(d_2(u)-1)(d_2(v)-1)} \\ &= n\sqrt{[(2n-2-1)(2n-4-1)]} + n\sqrt{[(2n-2-1)(0-1)]} \\ &\quad + n\sqrt{[(2n-4-1)(0-1)]} + n\sqrt{[(2n-4-1)(2n-4-1)]} \\ &= n\sqrt{(2n-3)(2n-5)} + n\sqrt{3-2n} + n\sqrt{5-2n} + n(2n-5) + n(2n-5) \\ &= n \left[\sqrt{[(2n-3)(2n-5)]} + \sqrt{3-2n} + \sqrt{5-2n} + (2n-5) \right] \end{aligned}$$

and

$$\begin{aligned} \text{LRM}_2(G) &= \sum_{uv \in E} (d_2(u)-1)(d_2(v)-1) \\ &= n[(2n-2-1)(2n-4-1)] + n[(2n-2-1)(0-1)] + n[(2n-4-1)(0-1)] \\ &\quad + n[(2n-4-1)(2n-4-1)] \\ &= n[(2n-3)(2n-5)] + n[(2n-3)(-1)] + n[(2n-5)(-1)] + n(2n-5)^2 \\ &= n(8n^2-40n+48) \\ &= 8n(n^2-5n+6) \end{aligned}$$

Theorem

Let Sf_n be a sunflower graph with $3n + 1$ vertices, its leap reduced reciprocal Randic and leap reduced second Zagreb indices are given below:-

$$\begin{aligned} \text{LRRR}(G) &= n \left[\sqrt{2-3n} + \sqrt{3-3n} + \sqrt{5-3n} + \sqrt{[(3n-3)(3n-5)]} \right. \\ &\quad \left. + (3n-5) \right] \\ \text{LRM}_2(G) &= n(18n^2-63n+50) \end{aligned}$$

Proof: We compute the required result by using definitions 1 and 2 given in the previous section as follows:-

$$\begin{aligned}
\text{LRRR}(G) &= \sum_{uv \in G} \sqrt{(d_2(u)-1)(d_2(v)-1)} \\
&= n\sqrt{[(3n-1-1)(0-1)]} + n\sqrt{[(3n-2-1)(0-1)]} + n\sqrt{[(3n-4-1)(0-1)]} \\
&\quad + n\sqrt{[(3n-2-1)(3n-4-1)]} + n\sqrt{[(3n-4-1)(3n-4-1)]} \\
&= n\sqrt{2-3n} + n\sqrt{3-3n} + n\sqrt{5-3n} + n\sqrt{[(3n-3)(3n-5)]} + n(3n-5) \\
&= n\left[\sqrt{2-3n} + \sqrt{3-3n} + \sqrt{5-3n} + \sqrt{[(3n-3)(3n-5)]} + (3n-5)\right]
\end{aligned}$$

$$\begin{aligned}
\text{LRM}_2(G) &= \sum_{uv \in G} (d_2(u)-1)(d_2(v)-1) \\
&= n[(3n-1-1)(0-1)] + n[(3n-2-1)(0-1)] + n[(3n-4-1)(0-1)] \\
&\quad + n[(3n-2-1)(3n-4-1)] + n[(3n-4-1)(3n-4-1)] \\
&= n(2-3n) + n(3-3n) + n(5-3n) + n[(3n-3)(3n-5)] + n(3n-5)^2 \\
&= n(18n^2-63n+50)
\end{aligned}$$

4. Conclusion

In this paper, we have extended the work initiated by Naji et al. In this regard, we have defined the leap reduced reciprocal Randic and leap reduced second Zagreb indices and we have computed the exact values of these newly defined indices for selected graphs including wheel graph, gear graph, helm graph, flower graph and sunflower graph.

References

- [1] Gutman I, Trinajstić N. Graph theory and molecular orbitals: total-electron energy of Alternant hydrocarbons. *Chem Phys Lett.* 1972;17:535–538.
- [2] Borovicanin KC, Das BF, Gutman I. Bounds for Zagreb indices. *MATCH Commun Math Comput Chem.* 2017;78(1):17–100.
- [3] Das KC, Gutman I. Some properties of the second Zagreb index. *MATCH Commun Math Comput Chem.* 2004;52:103–112.
- [4] Gutman I, Ruscic B, Trinajstić N, Wilcox CF. Graph theory and molecular orbitals XII acyclic polyenes. *J Chem Phys.* 1975;62(9):3399–3405.
- [5] Nikolic S, Kovacevic G, Milicevic A, Trinajstić N. The Zagreb indices 30 years after, Croat. *Chem Acta.* 2003;76:113–124.
- [6] Xu K, Hua H. A unified approach to extremal multiplicative Zagreb indices for trees, unicyclic and bicyclic graphs. *MATCH Commun Math Comput Chem.* 2012;68:241–256.

- [7] Gutman I. Multiplicative Zagreb indices of trees. *Bull Soc Math, Banja Luka*. 2011;18: 17–23.
- [8] Todeschini R, Consonni V. New local vertex invariants and molecular descriptors based on functions of the vertex degrees. *MATCH Commun Math Comput Chem*. 2010;64:359–372.
- [9] Ashra AR, Doslic T, Hamzeh A. The Zagreb coindices of graph operations. *Discrete Appl Math*. 2010;158:1571–1578.
- [10] Gutman I, Furtula B, Vukicevic ZK, Popivoda G. On Zagreb indices and coincides. *MATCH Commun Math Comput Chem*. 2015;74:5–16.
- [11] Xu K, Das KC, Tang K. On the multiplicative Zagreb coindex of graphs. *Opuscula Math*. 2013;33:197–210.
- [12] Eliasi M, Iranmanesh A, Gutman I. Multiplicative versions of first Zagreb index. *MATCH Commun Math Comput Chem*. 2012;68:217–230.
- [13] Naji AM, Soner ND, Gutman I. On leap Zagreb indices of graphs. *Communi Combin Optim*. 2017;2(2):99–117.
- [14] Naji AM, Soner ND. The first leap Zagreb index of some graph operations. *Int J Appl Graph Theory*. 2018;2(1):07–18.
- [15] Shiladhar P, Naji AM, Soner ND. Leap Zagreb indices of some wheel related graphs. *J Comput Math Sci*. 2018;9(3):221–23.
- [16] Dayan F, Javaid M, ur Rehman MA. On leap gourava indices of some wheel related graphs. *Sci Inquiry Rev*. 2018;2(4):14– 24.
- [17] Dayan F, Ahmad B, Zulqarnain M, Ali U, Ahmad Y, Zia TJ. On some topological indices of triangular silicate and triangular oxide networks. *Int J Pharm Sci Res*. 2018; 9(10): 4326–31. doi: 10.13040/IJPSR.0975-8232.9(10).
- [18] Dayan F, Javaid M, Zulqarnain M, Ali MT, Ahmad B. Computing banhatti indices of hexagonal, honeycomb and derived networks. *Am J Math Comput Modell*. 2018;3(2):38–45. doi: 10.11648/j.ajmcm.20180302.11
- [19] Dayan F, Javaid M, Ali U, Ahmad B, Zulqarnain M. On some banhatti indices of triangular silicate, triangular oxide, rhombus silicate and rhombus oxide networks. *Am J Inf Sci Technol*. 2018;2(2):42–49. doi: 10.11648/j.ajist.20180202.13